


Every day, more than 130 people in the United States die after overdosing on opioids.

In 2017, an estimated 1.7 million people in the U.S. suffered from substance use disorders related to prescription opioid pain relievers and 652,000 suffered from a heroin use disorder.



80%
of people who use heroin first misused prescription opioids.



21-29%
of patients prescribed opioids for chronic pain misuse them.




8-12%
develop an opioid use disorder.



Opioid overdoses increased 30 percent from July 2016 through September 2017 in 52 areas in 45 states.



The CDC estimates the total economic burden of prescription opioid misuse in the U.S. is **\$78.5 billion** per year.



An estimated 4 to 6 percent who misuse prescription opioids transition to heroin.



Opioid overdoses in large cities increase by 54 percent in 16 states.



Medication-Assisted Treatment




In-House Counseling Services



Personalized Support Services/Case Management

EVIDENCE OF EFFECTIVENESS



For most people, the use of medications combined with psychosocial treatment is superior to drug or psychosocial treatment on its own.
American Society of Addiction Medicine, The ASAM National Practice Guideline.



“Studies have shown that outcomes are much better when you are on medication-assisted therapy. For one, it decreases risk of relapse — significantly. Second, MAT has also been shown to be effective in preventing infectious diseases like HIV. Third, medication-assisted therapy has been shown to be effective in preventing overdoses.”

- Dr. Nora Volkow, the Director of the National Institute on Drug Abuse.

SAVIDA HEALTH OUTCOMES



60%

In the 1st month of treatment, 60% of our patients test negative for opiates.



91%

After 3 months, 91% of our patients test negative for opiates.



75%

In months 2 and 3 of treatment 75% of our patients test negative for opiates.



2 months of residential treatment costs as much as four years of treatment with Savida Health

(1) Based on patients' toxicology results in December 2018 and January 2019. SaVida is exploring methods to better track discharge rates to many with the toxicology data.

Good outcomes are contingent on adequate treatment length.

Generally, for residential or outpatient treatment, participation for less than 90 days is of limited effectiveness, and treatment lasting significantly longer is recommended for maintaining positive outcomes.